

## Psychologisches Institut – Psychopathologie und Klinische Intervention (Universität Zürich)

Í rafrænum meðferðum á geðdeild Háskólans í Zurich er lögð áhersla að að vinna með áföll, áfallastreituröskun, alvarleg sorgarviðbrögð og aðlögunarvandamál. Aðferðin miðar að því að draga úr persónulegum kvörtunum einstaklinganna og bæta lífsgæði þeirra. Meðferðin samanstendur af 12 til 16 fundum sem fara fram tvisvar í viku. Byggt er á hugrænni atferlismeðferð sem miðar að því að styrkja einstaklinga og fá þá til að endurskoða hugsanir sínar, viðurkenna truflun á hegðun og finna leiðir til að takast á við erfiðar aðstæður í lífinu.

Frekari leitarorð: Aðskilnaðarkvíði, Áfallastreita, Alvarleg sorg

Markhópur	Efni	Tegund	Land
• Fullorðnir	• Þunglyndi • Áfall	• Meðferð	• Sviss

The psychotherapeutic centre of the University of Zürich provides web-based therapies in the fields of trauma and post-traumatic stress disorders (PTSD), depression, persistent severe grief and adaptive disorders. The client is accompanied by a personal psychologist during the whole treatment. The therapy takes place via a secured internet platform and consists of 12 to 16 sessions. Each phase starts with a detailed introduction which explains the meaning of the practices. The client works on the tasks individually on the dates determined by him-or herself. The therapist reacts to the tasks within one work day. The basis of the modules is cognitive behavioural therapy which is scientifically proven to be appropriate for web-based therapies. The aim is to encourage the patient to question his/her thoughts, to recognize dysfunctional behaviours and to work out better strategies to cope with difficult life situations. The therapy for patients with trauma or post-traumatic stress disorder addresses people who had a traumatic experience at least three months ago and from which they still suffer. It consists of three phases which are self-confrontation, cognitive restructuring and sharing of fate. The therapy of depression includes five phases of treatment, these are: consideration of situations in which the patient is depressive, building up positive activities, become aware and change of negative and automatic thoughts, difficulties in interpersonal relationships as well as prevention of relapses. The treatment of persistent severe grief addresses people who sustained a traumatic loss and still suffer from that after six or more months. This kind of therapy is based on the same steps as the therapy of post-traumatic stress disorder. The treatments are not appropriate for people with addiction, dissociations, psychosis and suicidal tendencies.



## Upphaflegt tungumál

German

## Land

Switzerland

## **Tengill**

http://www.psychologie.uzh.ch/de/fachrichtungen/psypath/Psychotherapie/Onlintherapie.html

(access date: 07/04/2017)



CC - Attribution-NonCommercial-ShareAlike http://creativecommons.org/licenses/by-nc-sa/3.0/

ILI. *Psychologisches Institut – Psychopathologie und Klinische Intervention (Universität Zürich).* Retrieved on Monday August 25, 2025 from https://www.ecounselling4youth.eu/online-material/courses/TGP541/

Google Suche

Home | Kontakt | Sitemap & | RE

Framkvæmdastjórn Evrópusambandsins, styrkir verkefnið en er ekki ábyrgt fyrir skoðunum eða innihaldi efnis tengdu verkefninu.

https://www.ecounselling4youth.eu/project/